



Japan
Food
Research
Laboratories

Japan Food Research Laboratories

Authorized by the Japanese Government

HEAD OFFICE : 52-1 Motoyoyogi-cho, Shibuya-ku, Tokyo 151-0062
OSAKA BRANCH : 3-1 Toyotsu-cho, Suita-shi, Osaka 564-0051
NAGOYA BRANCH : 5-13 Oosu 4-chome, Naka-ku, Nagoya 460-0011
KYUSHU BRANCH : 1-12 Shimogofuku-machi, Hakata-ku, Fukuoka 812-0034
TAMA LABORATORY : 11-10 Nagayama 6-chome, Tama-shi, Tokyo 206-0025
CHITOSE LABORATORY : 2-3 Bunkyo, Chitose-shi, Hokkaido 066-0052

REPORT

No. 207062138-004 1/6

August 02, 2007

SAGAN COAT Photocatalyst Coating Solution TPX for Primary Skin Irritation in Rabbits

Requested by: Kon Corporation
22646 Matsunokihara, Yamauchi-cho, Takeo-city
Saga-prefecture 849-2305
Japan

Received: June 26, 2007

I, the undersigned, hereby declare that the work described in this report was performed under my supervision, as Study Director, in accordance with OECD Guidelines for the Testing of Chemicals 404 (2002), and that the report provides a true and accurate record of the results obtained.

This is a translation of the original report, No. 207062138-001, written in Japanese.

Study Director

Tomoko Shimazaki

Tomoko Shimazaki
Section of Biological Safety Research
Tama Laboratory
Japan Food Research Laboratories

August 15, 2007

Date



Other contributors:

Takeshi Nagai D.V.M.
Yasuharu Kawamoto
Mirai Ozawa
Misora Suzuki

SAGAN COAT Photocatalyst Coating Solution TPX for Primary Skin Irritation in Rabbits

1. Abstract

The test sample, SAGAN COAT Photocatalyst Coating Solution TPX, was tested for primary skin irritation in rabbits in accordance with OECD Guidelines for the Testing of Chemicals 404 (2002).

The test sample was applied occlusively for four hours to the intact and abraded skin of three rabbits. As a result, very slight to well-defined erythema was found in all of the animals at one hour after removal, but all the erythema disappeared at the 24-hour observation.

The primary irritation index (P.I.I.) calculated according to Federal Register (1972) was 0.4. Consequently, the category of the P.I.I. of the test sample was regarded as "Negligible".

2. Test sample

SAGAN COAT Photocatalyst Coating Solution TPX
Character: Yellowish suspended solution

3. Test period

From July 09 to August 02, 2007

4. Experimental animals

Three male Japanese white rabbits were purchased from Kitayama Labes Co., Ltd. They were acclimated to the laboratory conditions for more than one week to confirm that there were no abnormalities in general condition. They were individually housed in FRP cages under the standard laboratory conditions (Temperature: $22\text{ }^{\circ}\text{C} \pm 2\text{ }^{\circ}\text{C}$, Light-dark cycle: 12/12 hours). And they were given tap water *ad libitum* and a certain amount of LRC4 diet [Oriental Yeast Co., Ltd.] in proportion to each animal's body weight.

5. Procedures

The back of each animal was clipped and freed of hair about 24 hours before the test, and the shaved area was divided into four sections (approximately 6 cm²). Two of the test sections were abraded and the other two were left intact. Abrasion was made by epidermal incision through the stratum corneum; two incisions perpendicular to the other two were made in a manner resembling a “#” pattern by use of a sterile 18-gage needle.

A 0.5-mL portion of the test sample was applied to an approximately 2 cm × 3 cm gauze patch, which was held in one intact and one abraded sections on each animal with adhesive plaster [NICHIBAN Co., Ltd.]. To ensure the contact of the test sample with skin, the patches were covered with Blenderm surgical tape [3M Health Care Ltd.]. The other intact and the abraded sections served as control.

The exposure duration was four hours. After exposure, the patches were removed. All of the test sections were wiped with pure water. Readings of the skin reactions were made at 1, 24, 48 and 72 hours after removal of the patches based on Table 1.

According to Federal Register (1972), the primary irritation index (P.I.I.) was obtained as follows: For each animal, the scores of the readings at 1, 24 and 48 hours were added together, and the summation was divided by 6. The mean value for the three animals was calculated to obtain the P.I.I. The degree of primary skin irritation of the test sample was determined by means of P.I.I. on the criteria of ISO 10993-10 shown in Table 2.

The animals were weighed at the start and completion of the test.

6. Results (Tables 3 and 4)

Well-defined erythema (score 2) was found at intact skin of one animal (Rabbit No. 3) and very slight erythema (score 1) was found in the other skin at one hour after removal, but all the erythema was disappeared at the 24-hour observation, and no other abnormalities were found during the follow period.

The P.I.I., calculated by the scores of the readings, was 0.4.

At non-treatment sections, no irritation response was observed throughout the observation period.

7. Discussion

The test sample was tested for primary skin irritation in rabbits in accordance with OECD Guidelines for the Testing of Chemicals 404 (2002). As a result, very slight to well-defined erythema was found in all of the animals at one hour after removal, but all the erythema disappeared at the 24-hour observation.

The primary irritation index (P.I.I.), calculated according to Federal Register (1972), of the test sample was 0.4. Consequently, the category of the primary irritation index of the test sample was regarded as “Negligible”.

8. References

- Federal Register (§ 191, December, 1972).
- ISO 10993-10 Biological Evaluation of Medical Devices-Part 10: Tests for irritation and delayed-type hypersensitivity 6.3 Animal skin irritation test (2002).

Table 1. Evaluation of skin reactions

Erythema and eschar formation

No erythema	0
Very slight erythema (barely perceptible)	1
Well-defined erythema	2
Moderate to severe erythema	3
Severe erythema (beet redness) to slight eschar formation (injuries in depth)	4*
	[Total possible erythema score 4]

- * Bleeding, ulcer and necrosis, as considered injuries in depth, are classified as score 4.

Edema formation

No edema	0
Very slight edema (barely perceptible)	1
Slight edema (edges of area well-defined by definite raising)	2
Moderate edema (raised approximately 1 mm)	3
Severe edema (raised more than 1 mm and extending beyond area of exposure)	4
	[Total possible edema score 4]

Table 2. Irritation response categories in rabbit

Response category	P.I.I.
Negligible	0 to 0.4
Slight	0.5 to 1.9
Moderate	2 to 4.9
Severe	5 to 8

Table 3. Body-weight changes (kg)

Rabbit No.	Before treatment	Completion of the test
1	3.04	3.02
2	2.98	3.07
3	3.25	3.43

Table 4. Scores of irritation response

Observation time (hour)	Rabbit No. 1		Rabbit No. 2		Rabbit No. 3	
	Intact	Abraded	Intact	Abraded	Intact	Abraded
1	1/0	1/0	1/0	1/0	2/0	1/0
24	0/0	0/0	0/0	0/0	0/0	0/0
48	0/0	0/0	0/0	0/0	0/0	0/0
72	0/0	0/0	0/0	0/0	0/0	0/0

Values are expressed as scores of Erythema·Eschar/Edema.